

Effect of soil biota on growth and allocation by *Eucalyptus microcarpa*

Mark Bourne · Adrienne B. Nicotra ·
Matthew J. Colloff · Saul A. Cunningham

Received: 10 October 2007 / Accepted: 19 December 2007
© Springer Science + Business Media B.V. 2007

Abstract We examined growth of *Eucalyptus microcarpa* seedlings in soil collected from four sites in southeastern Australia, in which retired pasture land has been revegetated with mixed plantings of *Eucalyptus* and *Acacia* species. Revegetation of farm land in southeastern Australia is an area of major investment. The focus of the study was to examine the influence of soil biota on seedling growth and its possible interaction with soil enrichment from a legume (*Acacia*) and decomposition rates. We used a soil freezing treatment (-80°C for 3 days) to retard the soil biota, with the expectation that invertebrates in particular would be killed. Soil freezing did not cause a nutrient pulse, but did reduce the level of ammonium in soil. Nitrate levels increased with time in pots, regardless of the soil treatment. Decomposition rates measured using cellulose substrate were significantly reduced by the freeze treatment, but only for approximately 90 days. *Eucalyptus microcarpa* seedlings grown in freeze-treated soil were approxi-

mately 40% smaller (total biomass), had marginally lower LAR (leaf area ratio), and significantly lower LMA (leaf mass per area). Low LMA indicates that leaves are either thinner in cross-section or less dense. We hypothesise that both the poor growth of seedlings and production of less robust leaves are consequences of reduced availability of soil nutrients due to the diminished soil biota after freeze treatment. Litter under *Acacia* was richer in nitrogen than litter under *Eucalyptus* but there was no difference in nitrogen content of soil, and consequently no soil source effects on plant growth or decomposition. We suggest that variation in the soil biota has the potential to greatly enhance or hinder the success of revegetation on retired agricultural land, but enrichment of soil by decomposition of nitrogen rich litter in these sites requires longer than the 8–15 years since they were revegetated.

Keywords Allocation · Growth · Leaf · Legume · Nitrogen · Revegetation

Responsible Editor: Barbara Wick.

M. Bourne · A. B. Nicotra
School of Botany and Zoology,
Australian National University,
Canberra, ACT 0200, Australia

M. Bourne · M. J. Colloff · S. A. Cunningham (✉)
CSIRO Entomology,
GPO Box 1700, Canberra, ACT 2601, Australia
e-mail: saul.cunningham@csiro.au

Introduction

In the last decade or so, it has become widely recognised that soil biota play an important role in plant growth (Bonkowski et al. 2000; Poveda et al. 2005) and community composition (Bever et al. 1997; Bradford et al. 2002; De Deyn et al. 2004). There is now a greater appreciation of the inter-dependence of

aboveground and belowground subsystems and the interactions between soil biota and plants (Bardgett 2004; Wardle et al. 2004). The primary mechanisms underpinning the effects of soil biota on plant growth are the mineralisation of plant residues by decomposer organisms and the release of nutrients into the soil in plant-available form, and the capacity of the soil biota to directly assist plants in nutrient uptake. These processes are influenced by soil microbial biomass and composition, which is, in turn, dependent on the amount and quality of inputs of plant litter (Wardle 2002; Rantalainen et al. 2004).

The invertebrate component of the decomposer biota affects plant growth and resource allocation via indirect processes including: (1) alteration of microbial abundance and community composition (Klironomos et al. 1992, but see McLean et al. 1996); (2) enhancement of nutrient mineralisation and transfer (Haimi et al. 1992; Schmidt and Curry 1999; Baker et al. 2003); (3) influencing competition between plants (Kreuzer et al. 2004; Endlweber and Scheu 2006); (4) redistribution of organic matter and nutrients (Cortez and Bouché 1998) and (5) enhancing leaf herbivory (Scheu et al. 1999; Poveda et al. 2005). Furthermore, root herbivore invertebrates can influence plant growth and resource allocation both directly (Wardle 2002) and via indirect mechanisms such as the release of organic compounds via plant wounds and faecal pellets that subsequently become available for plants (Yeates et al. 1999).

The majority of research on the effects of soil biota on plant growth and community structure has been undertaken in Europe and North America, but Australian environments are often quite different. Australian soils tend to be old, strongly leached and nutrient-poor (McKenzie et al. 2004). The Australian flora is notable for the abundance of sclerophyllous species, which produce leaf litter that is tough (Read et al. 2005), and rich in plant secondary metabolites (Cooper 2001). The decomposer biota may be highly seasonally variable in diversity and abundance, because they are limited by soil moisture (Spain and Hutson 1983). Decomposition in the drier areas tends to be sporadic, contingent upon rainfall events (Wood 1974; Hutson and Veitch 1985) and fire (Radho-Toly et al. 2001). There are, therefore, strong reasons to suspect that northern hemisphere studies provide a poor basis for understanding aboveground–belowground interactions in Australia.

The genera *Acacia* and *Eucalyptus* dominate the Australian flora, often co-occurring. They each have mechanisms to cope with drought, fire and low levels of soil nutrients, especially nitrogen and phosphorus (Adams and Attiwill 1984; Gill 1994). The *Acacia* are legumes, often pioneer species, with symbiotic, rhizobial N-fixing capacity (Stock et al. 1995; Thrall et al. 2000), while *Eucalyptus* have mutualisms with mycorrhiza whereby they acquire N and P (Launonen et al. 2004). *Eucalyptus* are noted for their sclerophyllous foliage (another adaptation to nutrient poor soils) but many *Acacia* species produce less sclerophyllous leaves (or phyllodes) with higher nitrogen content. In this way the two genera usually differ in the quality of leaf litter they drop. The nitrogen economy (Wright et al. 2004) is also expected to greatly influence growth responses to environmental variation, potentially affecting both the rate of leaf production (with positive feedback into future growth) and the structure of the leaves produced.

Broad-scale revegetation, including natural regeneration, environmental plantings and agroforestry, is a major activity in the restoration and rehabilitation of degraded Australian landscapes (Dorrough and Moxham 2005; Vesik and Mac Nally 2006), involving revegetation targets of several millions of hectares nationwide, at a cost roughly in the order of Aust \$2,000–6,000 dollars per hectare (McCarthy and Lindenmayer 2007). *Eucalyptus* spp. are commonly the dominant element of revegetated sites and *Acacia* spp. are commonly co-planted in order to provide a source of nitrogen (Thrall et al. 2005). Many studies of mixed *Eucalyptus* and *Acacia* plantations have demonstrated superior growth compared to monocultures, but the outcomes are context dependent (Forrester et al. 2006). The interactions between plants and soil biota in relation to growth and resource allocation are therefore of considerable environmental and economic significance in relation to ecological restoration in Australia.

One approach to assess the overall role of soil organisms as drivers of plant response is to examine the effect of soil sterilisation. Various methods have been used in order to retard populations of the soil biota. Selective chemical biocides for invertebrates, fungi and bacteria have been used to examine the effects of these taxa on decomposition and mineralisation in the field (Beare et al. 1992; Hu et al. 1995).

Others have used gamma-irradiation (Troelstra et al. 2001). For this experiment we chose to use soil freezing because it posed the least risk of introducing direct effects on soil nutrient status (such as associated with chemical intervention) and because it is expected to have a differential effect on invertebrates. Freezing at the conditions we used is likely to kill invertebrates outright (Bardgett and Chan 1999) while spores of fungi and bacteria are likely to remain viable (Vishnivetskaya et al. 2000; Walker et al. 2006).

We used pots containing soil and soil biota from revegetation sites established on retired pasture land in order to measure decomposition rates and the growth responses of Grey Box, *Eucalyptus microcarpa*, a common tree species in south-eastern Australia. We compared effects of a soil freezing treatment and a nitrogen supply treatment in the form of soil taken from under the legume, *Acacia pycnantha*.

The purpose of the study was to ask:

- (1) Do soil biota affect growth of *E. microcarpa* seedlings, as well as the allocation of resources to different parts of the plant?
- (2) Does soil sourced from under a legume (*Acacia pycnantha*) provide more nitrogen for growth of *E. microcarpa* seedlings, and is availability of this nitrogen influenced by the soil freezing treatment?
- (3) Does the soil source or soil freezing treatment influence decomposition rates in soil?

Materials and methods

Sites, soil sampling and construction of pots

Soil and litter was collected from sites located in north-central Victoria, Australia; near Charlton (36° 25'S, 143°27' E), Bendigo (36°54'S, 144°00'E), Heathcote (37°02'S, 144°43'E) and St Arnaud (36° 51'S, 143°04'E) between 12 April and 14 April 2005. These were revegetation sites on retired grazing land, planted with a mixture dominated by *Eucalyptus* and *Acacia* species. All sites had both grey box, *Eucalyptus microcarpa* and golden wattle, *Acacia pycnantha*. Site characteristics were (in the following order: Charlton, Bendigo, Heathcote, St Arnaud), year planted: 1997, 1994, 1996, 1990; range of diameter (cm at breast height) of *E. microcarpa* individuals

from under which soil was sampled: 6–10, 12–17, 8–15, 4–17; mean distance (m) between canopies: 2, 0, 4, 1.

At each site, soil and litter were collected beneath four each of *A. pycnantha* and *E. microcarpa* trees. We selected an area of 0.25 m² that represented the most complete litter cover within 2 m of the base of each tree, and then collected separate fractions of leaf litter, shallow soil (0–5 cm) and deep soil (5–15 cm). Each fraction was sealed in a plastic bag and transported to the laboratory in insulated coolers.

Soil and litter layers were reassembled in pots (14 cm diameter×15 cm deep; 2.2 l) within 5 days of collection. Deep soil was placed into pots to a depth of 10 cm, followed by 5 cm of shallow soil, followed by approximately 40 g of litter. The soil and litter sampled from under each tree was used to construct one pot for freeze-treatment, one control, and contributed half the material for the mixed pot (Fig. 1). For the mixed pots we combined soil, 1:1, from one *Eucalyptus* tree with soil from one *Acacia* tree. This latter treatment was included to help distinguish between two possible mechanisms associated with an *Acacia* soil benefit. If soil under *Acacia* has more plant-available nitrogen, one would predict that the plant response would be dosage dependent, with seedlings in mixed pots intermediate between those in the 'pure' soil treatments. Alternatively, *Acacia* soil might be beneficial because it is free of *Eucalyptus* specific organisms that harm growth (e.g. root feeding nematodes). In this case one would predict that the mixing of soils inoculates pots with harmful organisms, so that seedling performance in mixed pots would be more like that seen in pure *Eucalyptus* soil than in pure *Acacia* soil.

Experimental design

Soil treatment (freezing vs. control) and soil source (soil from under *E. microcarpa* or *A. pycnantha* or mixed) were applied orthogonally, creating 6 unique combinations. With four replicates of each, there were 24 pots in each block. Pots were arranged on benches in a greenhouse in a randomised block design (Fig. 1). Combining site with greenhouse block allowed for a strong test of the treatments of interest, but limited our capacity to distinguish between site effects (from the field) and block effects (from the greenhouse). Blocks were rotated clockwise within the greenhouse fort-




Fig. 1 Study design for greenhouse pot trial. Blocks refer to the collecting sites and also the layout of pots in the greenhouse. Source refers to the tree species from under which soil and litter were collected, and treatment refers to freezing versus controls. Numbers 1–4 refer to individual trees at each site, and indicate the distribution into pots of soil and litter that was sampled from under each tree, e.g. 1,1,=half the soil in the

mixed pot came from *A. pycnantha* tree 1 and half from *E. microcarpa* tree 1. *Shaded cells*, soil in pots that were planted with *E. microcarpa* seedlings. *Unshaded cells*, pot not planted with seedling but was used for monitoring of soil nitrogen. Only the first block is illustrated in detail, as all other blocks follow the same model

nightly to minimise positional effects. Contamination of pots by crawling invertebrates was minimised by applying fluon to the rim of the pots.

Soil freezing treatment and planting

To retard biological activity in the soil, particularly of invertebrate fauna, half the pots were freeze-treated before planting by holding assembled pots of soil and litter at -80°C for 3 days. Pots were subsequently kept in a greenhouse where the temperature ranged from 5 to 33°C during the course of the experiment (April to September). Pots were kept well watered before seedlings were planted.

Eucalyptus microcarpa seeds were supplied by the National Tree Seed Centre (ENSIS, Canberra). To minimise genetic variation we used seed collected from the same mother tree on the same date. Seeds were germinated in 1:1 vermiculite/perlite and then provided with liquid fertiliser for about 2 weeks prior to planting. Seedlings were planted in pots when they reached a height of 5 mm. Germination and early growth took longer than anticipated, so pots were left in the greenhouse for some time after the soil freezing

treatment and before seedlings were planted. After planting, pots were watered close to saturation twice weekly. We judged that pots were saturated by feeling the weight of pots, and observing drainage.

Three blocks were planted 23 days after freezing treatment (17 May 2005) and the fourth block was planted 7 days later (24 May 2005). Sixteen pots with soil and litter from the mixed source were not planted with a seedling and were only used to detect any post-freezing nutrient pulse (discussed further below; Fig. 1).

Estimation of decomposition rates

We used filter paper discs (Whatman, 42.5 mm diameter) as a standard cellulose substrate to assess decomposition rates (Deacon 1985; Savoie and Goubière 1989; Carter et al. 1999). Discs were placed beneath the litter layer 33 days after freezing treatment, shortly after the seedlings were planted. All 80 pots containing seedlings were used. Discs were left in situ until those that decomposed most rapidly had almost disappeared (27 May–29 June). The remains were removed, dried (70°C for 24 h) and

weighed to determine mass loss. Observations were repeated, using fresh filter papers, between 29 June and 27 July (27 days) and 1–22 August (21 days).

Measurements of plant growth and biomass allocation

Plants were destructively sampled on 6th and 7th September, 106 days after planting. Plants in the fourth block grew for 1 week less (99 days) than plants in the other three blocks, which contributes another source of variation to the block effect. Plants were removed from pots and then separated into stems, leaves and roots. Soil was tipped out of the pots and root fragments retrieved for a standard 2 min search time per pot. Plant parts were dried at 60°C for 72 h before weighing. Leaf area was measured using a flatbed scanner prior to drying.

Over the course of the experiment four plants died from unknown causes (evenly among the treatments, with no more than one from any of the six categories), reducing the overall sample size to 76 plants available for harvest. Six plants (three each from control and freeze treated pots) did not produce enough leaf tissue to perform nitrogen analyses, therefore only 70 seedlings were measured for leaf nitrogen content.

Carbon and nitrogen analysis

To test if there was a soil nutrient pulse after freezing, a sample of approximately 10 g of shallow soil was taken during weeks 1, 2 and 4 following the treatment, from 16 pots containing mixed soil without any seedling planted in them. Mineral nitrogen from soil samples was extracted in 2 M KCl (Rayment and Higginson 1992) and analysed colorimetrically for nitrate and ammonium using an Alpkem continuous flow analyser.

Total carbon and total nitrogen content of shallow soil, leaf litter and leaves of *E. microcarpa* seedlings was analysed according to the method of Dumas (1981), using a mass spectrometer (Europa 20–20) with an automated nitrogen and carbon analysis preparation system. Repeated N analyses on 11 leaf samples and 24 litter and soil samples were used to assess accuracy. Regression of repetition 1 versus repetition 2 of these samples explained 99.3% for leaf N and 99.8% of variation for litter/soil N measurements respectively. Coefficient of variation for %N measurement of standards was 0.68% for leaves ($n=$

42), 0.94% for litter ($n=27$) and 4.19% for soil ($n=27$). Together these assessments indicate a high level of accuracy in N analysis.

Data analysis

All data were analysed using GenStat (Lawes Agricultural Trust 2002). Data were log-transformed when necessary to meet the assumption of normal distribution of residuals. Where sample sizes were uneven (decomposition and plant allocation measurements) we used restricted estimates maximum likelihood (ReML) models with Wald significance tests (Payne 2005).

Freezing treatment and soil source were fixed factors and block (combining field site and greenhouse location) was a random factor for the analysis of plant growth, allocation, leaf traits and decomposition rates. Interaction terms were dropped if no significant interaction was found, but non-significant treatment effects were maintained. To analyse the effect of species (*E. microcarpa* vs *A. pycnantha*) on C and N content of leaves, litter and shallow soil material we used one-way analysis of variance (ANOVA) with species as a fixed factor and site as a random factor. To analyse the effect of freezing treatment on NO_3 , NH_4 and total mineral nitrogen ($\text{NO}_3 + \text{NH}_4$) in soil, we used both one-way ANOVA and repeated measures ANOVA.

Results

Soil carbon and nitrogen and plant growth

Litter and soil used to construct pots varied in carbon and nitrogen content depending on its source. Litter beneath *A. pycnantha* was significantly higher in nitrogen ($F_{1,27}=39.11$, $P<0.001$; Fig. 2) and had a lower C/N ratio compared with litter beneath *E. microcarpa* ($F_{1,27}=23.13$, $P<0.001$; data not shown). Soil from under *A. pycnantha* had a significantly lower C/N ratio than soil from under *E. microcarpa* ($F_{1,27}=12.04$, $P=0.002$), but the difference was small, and the nitrogen content was not significantly different ($F_{1,27}=1.61$, $P=0.22$; Fig. 2). In spite of these differences in quality between soil and litter from under *A. pycnantha* and *E. microcarpa*, there were no significant effects on plant growth, plant




Fig. 2 Mean total nitrogen content of litter and soil (\pm SE) comparing material collected beneath *A. pycnantha* and *E. microcarpa*

allocation or decomposition associated with the soil source, including the mixed treatment.

There was no evidence of a soil nutrient pulse following the freezing treatment (Fig. 3). There was a marginally significant decrease in ammonium following the freeze-treatment ($F_{1,11}=4.5$, $P=0.057$) and no significant effect of time ($F_{2,28}=1.2$, $P=0.314$) or interaction with time. By way of contrast, nitrate was not significantly affected by freezing, but increased significantly over the 4 week period (Fig. 3, $F_{2, 28}=23.16$, $P<0.001$) with no time-by-treatment interaction. The increase in nitrate content, combined with reduction in ammonium, suggests that nitrogen cycling activity recommenced in the soils relatively soon after freezing treatment.

Decomposition rates

Mass loss of filter paper loss ranged from 0 to 100% over all three observation periods. The pattern of loss varied from holes in otherwise complete paper discs to fragmentation of the paper. Increased fungal and bacterial activity was observed around areas of filter paper that had been decomposed. The fungal colonies present were consistent in appearance with *Rhizopus*, *Mucor*, *Fusarium* and *Stachybotrys* spp. (Celeste Linde, Australian National University, personal observations). Estimated mean daily loss of filter paper was greater in the control pots during the first 33 days




Fig. 3 Mean nitrate (NO₃) and ammonium (NH₄) content of soil in pots, 1, 2 and 3 weeks after freeze treatment. Values are means of \log_e -transformed data, back transformed for the figure. The standard errors were scaled to maintain the relationship to the mean after back transformation. Hatched columns show values for freeze-treated pots, open columns show values for control pots




Fig. 4 Daily decomposition rate, estimated as percentage loss of filter paper (mean \pm SE) following freezing treatment of soils, as measured over three separate observation events, using fresh filter paper on each occasion

(Wald=6.16, $P=0.013$; Fig. 4). During the second observation period (66–93 days post-freezing treatment) this difference was still marginally significant (Wald=2.74, $P=0.098$) but in the third period (99–120 days) there was no significant difference (Wald=0.75, $P=0.387$). Repeated measures ANOVA supported this analysis, showing a time-by-treatment interaction ($F_{2,177}=3.53$, $P=0.033$), and that daily loss of filter paper increased over the three observation periods ($F_{2,177}=30.89$, $P<0.001$). There was no effect of soil source on decomposition in any observation period.

Measurements of plant growth and allocation

Leaf, root and total biomass of *Eucalyptus microcarpa* seedlings grown in freeze-treated soil were each almost 40% lower than those grown in untreated soil (Table 1, Fig. 5a), although only leaf and total biomass reached statistical significance. There was no difference in stem mass between the treatments.

Leaf area and leaf area ratio (LAR; the ratio of leaf area to total plant weight) were lower in plants from freeze-treated soil compared with controls, but significance of this effect was marginal (Table 1). Leaf mass per leaf area (LMA, g/cm^2 ; Fig. 5b) was significantly lower in plants from freeze-treated soil (Table 1). In other words, *E. microcarpa* seedlings grown in freeze-treated soil produced smaller, flimsier leaves and less leaf tissue relative to other plant tissues. Other allocation variables were not significantly different between treatments (Table 1).

Nitrogen content per unit leaf mass showed a significant inverse correlation with LMA ($R^2=0.26$, $P<0.001$, $n=70$; Fig. 6). If N content per unit leaf area was completely independent of LMA then the relationship between LMA and N would have had a slope of -1 , i.e. a doubling of LMA leads to a halving of N per mass. However, the slope of LMA versus N per mass was in fact less than -1 , (Fig. 6) because of a countervailing positive correlation between LMA and N per unit leaf area ($R^2=0.21$, $P<0.001$, $n=70$; data not shown). Consequently, there was no significant effect of soil treatment on nitrogen content in leaves, such as one would have expected if LMA was the sole driver of nitrogen content. In other words, thinner (low LMA) leaves generally contained higher concentrations of nitrogen, but the slope and fit of this relationship was not great enough to lead to a




Fig. 5 Mean biomass and allocation to plant parts of *E. microcarpa* seedlings grown in freeze-treated and control pots. Values are means of \log_e -transformed data, back transformed for the figure. The standard errors were scaled to maintain the relationship to the mean after back transformation. **a** The total biomass of seedlings, and its allocation into leaves (dots), shoots (hatched) and roots (solid). The standard error applies to the total biomass. **b** Leaf mass area (LMA, g/m^2) of leaves from seedlings

significant effect of soil treatment on N per mass in leaves.

Discussion

The main effect of the soil freezing treatment on performance of *Eucalyptus microcarpa* seedlings was to significantly retard growth. This was evidenced by lower total and leaf biomass compared with controls, and the production of less robust leaves, as indicated by lower leaf mass per area (LMA). Together these effects suggest that the soil biota in untreated soils

Table 1 Effect of soil and litter treatment on growth and allocation of *E. microcarpa*

Variable	Wald statistic	P value
Leaf mass per area	6.41	0.011
Leaf mass	4.56	0.033
Total biomass	4.37	0.037
Leaf area	3.68	0.055
Root mass	3.43	0.064
Leaf area ratio	3.39	0.066
Stem mass	1.46	0.226
Root shoot ratio	0.06	0.800
Root mass ratio	0.06	0.809
Leaf mass ratio	0	0.965

Data were analysed using a ReML model, with $n=76$, 1 *df*, with soil treatment as a fixed effect, and block as a random effect. All data were \log_e -transformed before analysis.

was providing a net benefit to *E. microcarpa* seedlings, facilitating faster growth and development of more robust leaves.

Two lines of evidence support the hypothesis that the beneficial effect of the soil biotic community on plant growth is due to decomposition and nutrient cycling. The freezing treatment reduced the decomposing capacity of the soil, based on the filter paper bioassays, for at least 90 days, and reduced soil ammonium concentrations for at least 30 days. Higher rates of decomposition and nitrogen mineralisation in untreated soil may have been responsible for releasing more nitrogen in plant-available form. For *Eucalyptus*, ammonium availability appears to be particularly important in this regard. Several *Eucalyptus* spp. have been found to take up ammonium and the amino acid glycine considerably more rapidly and in greater amounts than nitrate (Warren 2006).

Plants have several mechanisms for avoiding rate-limited processes in soil nitrogen mineralisation. Where rates of nitrification are low, plants may take up ammonium in preference to nitrate. Where ammonification rates are low, plants may take up amino acids. Early successional species appear to have a higher capacity for nitrate uptake than species of later succession, which show preferential uptake of ammonium and amino acids (Chapin et al. 1993; Persson and Näsholm 2001). Ammonium and amino acids may be the dominant forms of nitrogen taken up by *Eucalyptus* spp. in Australian soils which can have quite high levels of ammonium and soluble organic

nitrogen relative to nitrate (Turnbull et al. 1996; Garnett and Smethurst 1999).

The freezing treatment could also have harmed the mycorrhizal associates of *E. microcarpa*. Mycorrhiza of *Eucalyptus* spp. are common and widespread in Australia, are best known in relation to phosphorus nutrition of their host plants (Burgess et al. 1993) and can markedly increase growth of *Eucalyptus* seedlings (Chen et al. 2006). Mycorrhiza can also mobilise sources of nitrogen otherwise unavailable to plants (Read and Perez-Moreno 2003). Turnbull et al. (1995) found that mycorrhizal seedlings of *Eucalyptus* spp. grew well on a range of organic sources of nitrogen, whereas non-mycorrhizal seedlings could only grow on mineral sources of nitrogen or glutamine, indicating that mycorrhiza confer on *Eucalyptus* spp. the capacity to use nitrogen from a broad range of sources.

The capacity of *Eucalyptus* to exploit a wide range of nitrogen sources is likely to provide flexibility in exploiting a range of low nutrient environments. However, it may also mean they are ecologically linked to a wider range of soil organisms than solely the bacterial populations involved in nitrite oxidation. For example, they may be more affected by heterotrophic invertebrate populations and the fungi involved in decomposition of peptides and their




Fig. 6 The relationship between leaf mass per area (*LMA*) and nitrogen content on a mass basis. The dashed line shows the relationship that would hold if N per area was held constant. The solid line shows the relationship (from linear regression) in the data collected, pooling treated and untreated pots because no significant difference in slope was attributable to the treatment effect

ammonification, as well as bacteria involved in the various steps in the nitrification pathway.

Bardgett et al. (1998) completely eliminated nematodes, microarthropods and larger fauna from soils by freezing soil samples at -80°C and air drying. Only spore-forming microorganisms are likely to withstand these conditions (Hughes and Nobbs 2004). Antarctic soil metazoa can survive -80°C for years (Newsham et al. 2006), but lethal temperatures of -5°C to -30°C are the norm for invertebrates from temperature latitudes (Brown and Gaugler 1996; Addo-Bediako et al. 2000). Our study showed that decomposition rate and nitrate concentration in pots was lowest immediately after the freezing treatment, but then increased with time. Both patterns suggest that microflora recolonised pots via airborne spores or increased from populations that remained in a resting stage after freezing. It is much less likely that any invertebrates (or their eggs and cysts) could have survived the prolonged extreme low temperatures, but they could reestablish by airborne dispersal into the greenhouse, or by crawling across the fluon barrier. Dipteran larvae were observed in freeze-treated pots towards the end of the experiment.

Litter under *A. pycnantha* was approximately 50% richer in nitrogen than litter under *E. microcarpa*. Nevertheless, there was no evidence of soil or litter source effects in this study on plant growth or decomposition rate. This lack of a nitrogen enrichment effect might in part be explained by the fact that the difference in soil nitrogen concentration was so small as to be non-significant. It might be that decomposition rates in the field are slow, so that most of the nitrogen remains relatively unavailable in the litter, with no significant enrichment of the soil over the short time that these sites have been under revegetation. Alternatively, available nitrogen may have been assimilated into the soil biota, but not subsequently released in a plant available form. Decomposition rates in the natural environment are likely to be extremely temporally variable, because of the great variability in rainfall, but we probably increased decomposition rates in pots, relative to the field, by frequent watering. The small soil mass in pots (relative to the field) could have lead to higher temperatures, also increasing decomposition.

Absence of a soil and litter source effect also indicates that there were not significant numbers of *E. microcarpa* specialist root feeders to create a 'home

site disadvantage'. Although root feeding has been important in some studies (De Deyn et al. 2004), we found that the abundance of plant feeding nematodes in our soil was relatively low (unpublished observations). Because there was no evidence of a home site disadvantage associated with soil from under *Eucalyptus*, or nitrogen enrichment of soil from under *Acacia*, one would expect that the mixed soil treatment would not be different to the pure soil treatments, and this was indeed the case.

Inoculation experiments, in which soil organisms are introduced into pots, have detected positive effects on plant growth associated with soil organisms (Alpei et al. 1996; Laakso and Setälä 1999; Liiri et al. 2002). Our study differs from these studies in focusing on a tree species rather than grasses or herbs, and in detecting an effect on leaf structure (i.e. LMA) as well as growth rates. Evolution of high LMA can be an adaptation to low nutrient and low rainfall environments, where selection favours robust long-lived leaves (Cunningham et al. 1999). However, when considering within-population variation in LMA, as in the present study, variation is more likely to reflect a developmental response. We suggest that low LMA in the plants grown in freeze-treated soils is a stress response to low nutrient availability. Low LMA leaves are less robust, and therefore less likely to remain functional for as long as high LMA leaves, so this response is unlikely to be adaptive. Leaf nitrogen analyses indicated that there was no difference in nitrogen concentration between treatments, so the LMA response does not appear to be primarily attributable to a nitrogen allocation strategy. There was, however, a correlation between LMA and N_{mass} and N_{area} , which was similar regardless of treatment, suggesting close coupling in these traits that may represent a constraint to variation in leaf structure.

Conclusions

This experiment demonstrates that the soil biota confers large benefits to the tree *Eucalyptus microcarpa*. Tree seedlings grown in freeze-treated soil not only suffered in terms of biomass, but also had less robust leaves. This effect of soil biota on leaf development underlines the potential for complex consequences of above-ground below-ground ecological linkages. Our data on the nitrogen dynamics and

decomposition rates support the hypothesis that the soil biota provided benefits through the release of nitrogen from complex compounds into plant-available forms. In this study we detected no legume enrichment of the soil, probably because insufficient time had passed since planting to allow decomposition of the nitrogen rich legume litter. We suggest that variation in the soil biotic communities has the potential to greatly enhance or hinder the success of revegetation on retired agricultural land. Benefits to the soil and soil fauna from legume enrichment are likely to occur, but may take more than a decade to be realised.

Acknowledgments We thank Alison Jeavons, Lindsay Ezard and Geoff Winsell (Department of Primary Industry, Victoria) for help with site selection, and the landholders for allowing us access to sites. Kim Pullen provided assistance in field. We thank Celeste Linde (Australian National University), Elizabeth Lindsay and Mike Hodda (CSIRO Entomology) for methodological advice and information on various groups of soil organisms. Financial support for this research was provided by the CSIRO Water for a Healthy Country Flagship.

References

- Adams MA, Attiwill PM (1984) Role of *Acacia* spp. in nutrient balance and cycling in regenerating *Eucalyptus regnans* F. Muell. forests. I. Temporal changes in biomass and nutrient content. *Aust J Bot* 32:205–215
- Addo-Bediako A, Chown SL, Gaston KJ (2000) Thermal tolerance, climatic variability and latitude. *Proc R Soc B* 267:387–410
- Alpei J, Bonkowski M, Scheu S (1996) Protozoa, nematoda and lumbricidae in the rhizosphere of *Hordelymus europaeus* (Poaceae): faunal interactions, response of microorganisms and effects on plant growth. *Oecologia* 106: 111–126
- Baker GH, Amato M, Ladd J (2003) Influences of *Aporrectodea trapezoides* and *A. rosea* (Lumbricidae) on the uptake of nitrogen and yield of oats (*Avena fatua*) and lupins (*Lupinus angustifolius*). *Pedobiologia* 47:857–862
- Bardgett R (2004) The biology of soil: a community and ecosystem approach. Oxford University Press, Oxford
- Bardgett RD, Chan KF (1999) Experimental evidence that soil fauna enhance nutrient mineralization and plant nutrient uptake in montane grassland ecosystems. *Soil Biol Biochem* 31:1007–1014
- Bardgett RD, Keiller S, Cook R, Gilburn AS (1998) Dynamic interactions between soil animals and microorganisms in upland grassland soils amended with sheep dung: a microcosm experiment. *Soil Biol Biochem* 30:531–539
- Beare MH, Parmelee RW, Hendrix PF, Cheng W, Coleman DC, Crossley DA (1992) Microbial and faunal interactions and effects on litter nitrogen and decomposition in agroecosystems. *Ecol Monogr* 62:569–591
- Bever JD, Westover KM, Antonovics J (1997) Incorporating the soil community into plant population dynamics: the utility of the feedback approach. *J Ecol* 85:561–573
- Bonkowski M, Cheng X, Griffiths BS, Alpei J, Scheu S (2000) Microbial–faunal interactions in the rhizosphere and effects on plant growth. *Europ J Soil Biol* 36:135–147
- Bradford MA, Jones TH, Bardgett RD, Black HI, Boag B, Bonkowski M, Cook R, Eggers T, Gange AC, Grayston SJ, Kandeler E, McCraig AE, Newington JE, Prosser JI, Setälä H, Staddon PL, Tordoff GM, Tscherko D, Lawton JH (2002) Impacts of soil faunal community composition on model grassland ecosystems. *Science* 298:615–618
- Brown IM, Gaugler R (1996) Cold tolerance of steinernematid and heterorhabditid nematodes. *J Thermal Biol* 21:115–121
- Burgess TI, Malajczuk N, Grove TS (1993) The ability of 16 ectomycorrhizal fungi to increase growth and phosphorus uptake of *Eucalyptus globulus* Labill. and *E. diversicolor* F. Muell. *Plant Soil* 153:155–164
- Carter JP, Spink J, Cannon PF, Daniels MJ, Osbourn AE (1999) Isolation, characterization, and avenacin sensitivity of a diverse collection of cereal-root-colonizing fungi. *Appl Environ Microbiol* 65:3364–3372
- Chapin FS, Moilanen L, Kielland K (1993) Preferential use of organic nitrogen for growth by a non-mycorrhizal arctic sedge. *Nature* 361:150–153
- Chen YL, Dell B, Malajczuk N (2006) Effect of *Scleroderma* spore density and age on mycorrhiza formation and growth of containerized *Eucalyptus globulus* and *E. urophylla* seedlings. *New Forests* 31:453–467
- Cooper PD (2001) What physiological processes permit insects to eat *Eucalyptus* leaves? *Austral Ecol* 26:556–562
- Cortez J, Bouché MB (1998) Field decomposition of leaf litters: earthworm–microorganism interactions—the ploughing-in effect. *Soil Biol Biochem* 30:795–804
- Cunningham SA, Summerhayes B, Westoby M (1999) Evolutionary divergences in leaf structure and chemistry, comparing rainfall and soil nutrient gradients. *Ecol Monogr* 69:569–588
- Deacon JW (1985) Decomposition of filter-paper cellulose by thermophilic fungi acting singly, in combination, and in sequence. *Trans Br Mycol Soc* 85:663–669
- De Deyn GB, Raaijmakers CE, van der Putten WH (2004) Plant community development is affected by nutrients and soil biota. *J Ecol* 92:824–834
- Dorrough J, Moxham C (2005) Eucalypt establishment in agricultural landscapes and implications for landscape-scale restoration. *Biol Conserv* 123:55–66
- Dumas JB (1981) Sur les procédés de l'analyse organique. *Annales de Chimie* 47:195–213
- Endlweber K, Scheu S (2006) Effects of Collembola on root properties of two competing ruderal plant species. *Soil Biol Biochem* 38:2025–2031
- Forrester DI, Bauhus J, Cowie AL, Vanclay JK (2006) Mixed-species plantations of *Eucalyptus* with nitrogen-fixing trees: a review. *For Ecol Manag* 233:211–230
- Garnett TP, Smethurst PJ (1999) Ammonium and nitrate uptake by *Eucalyptus nitens*: effects of pH and temperature. *Plant Soil* 214:133–140
- Gill AM (1994) Patterns and processes in open-forests of *Eucalyptus* in Southern Australia. In: Groves RH (ed)

- Australian vegetation. 2nd edn. Cambridge University Press, Cambridge, pp 197–226
- Haimi J, Huhta V, Boucelham M (1992) Growth increase of birch seedlings under the influence of earthworms: a laboratory study. *Soil Biol Biochem* 24:1525–1528
- Hu S, Coleman DC, Hendrix PF, Beare MH (1995) Biotic manipulation effects on soil carbohydrates and microbial biomass in a cultivated soil. *Soil Biol Biochem* 27:1127–1135
- Hughes KA, Nobbs SJ (2004) Long-term survival of human faecal microorganisms on the Antarctic Peninsula. *Antarctic Science* 16:293–297
- Hutson BR, Veitch LG (1985) Relationships between litterfall rate, litter mass and decomposition rate in *Eucalyptus* forests in southeastern Australia. *Aust J Ecol* 10:443–450
- Klironomos JN, Widden P, Deslandes I (1992) Feeding preferences of the collembolan *Folsomia candida* in relation to microfungus successions in decaying litter. *Soil Biol Biochem* 24:685–692
- Kreuzer K, Bonkowski M, Langel R, Scheu S (2004) Decomposer animals (Lumbricidae, Collembola) and organic matter distribution affect the performance of *Lolium perenne* (Poaceae) and *Trifolium repens* (Fabaceae). *Soil Biol Biochem* 36:2005–2011
- Laakso J, Setälä H (1999) Sensitivity of primary production to changes in the architecture of belowground food webs. *Oikos* 87:57–64
- Launonen TM, Ashton DH, Kelliher KJ, Keane PJ (2004) The growth and P acquisition of *Eucalyptus regnans* F. Muell. seedlings in air-dried and undried forest soils in relation to seedling age and ectomycorrhizal infection. *Plant Soil* 267:179–189
- Lawes Agricultural Trust (2002) GenStat, 6th edn. VSN International, Oxford, UK
- Liiri M, Setälä H, Haimi J, Pennanen T, Fritze H (2002) Relationship between soil microarthropod species diversity and plant growth does not change when the system is disturbed. *Oikos* 96:137–149
- McCarthy MA, Lindenmayer DB (2007) Info-gap decision theory for assessing the management of catchments for timber production and urban water supply. *Environ Manag* 39:553–562
- McKenzie N, Jacquier D, Isbell R, Brown K (2004) Australian soils and landscapes: an illustrated compendium. CSIRO Publishing, Melbourne
- McLean MA, Laneko N, Parkinson D (1996) Does selective grazing by mites and collembolan affect litter fungal community structure? *Pedobiologia* 40:97–105
- Newsham KK, Maslen NR, McInnes SJ (2006) Survival of Antarctic soil metazoans at –80 degrees C for six years. *CryoLett* 27:291–294
- Payne RW (2005) REML analysis of mixed models. In: Payne RW, Harding SA, Murray DA, Soutar DM, Baird DB, Welham SJ, Kane AF, Gilmour AR, Thompson R, Webster R, Wilson GT (eds) The guide to GenStat release 8, part 2: statistics. VSN International Ltd and Lawes Agricultural Trust (Rothamsted Experimental Station), Oxford, UK, pp 557–654
- Persson J, Näsholm T (2001) Amino acid uptake: a widespread ability among boreal forest plants. *Ecol Lett* 4:434–438
- Poveda K, Steffan-Dewenter I, Scheu S, Tschamtkte T (2005) Effects of decomposers and herbivores on plant performance and aboveground plant–insect interactions. *Oikos* 108:503–510
- Radho-Toly S, Majer JD, Yates C (2001) Impact of fire on leaf nutrients, arthropod fauna and herbivory of native and exotic eucalypts in Kings Park, Perth, Western Australia. *Austral Ecol* 26:500–506
- Rantalainen M-L, Kontiö L, Haimi J, Fritze H, Setälä H (2004) Influence of resource quality on the composition of soil decomposer community in fragmented and continuous habitat. *Soil Biol Biochem* 36:1983–1996
- Rayment GE, Higginson FR (1992) Australian laboratory handbook of soil and water chemical methods. Inkata Press, Port Melbourne
- Read DJ, Perez-Moreno J (2003) Mycorrhizas and nutrient cycling in ecosystems—a journey towards relevance? *New Phytol* 157:475–492
- Read J, Sanson GD, Lamont BB (2005) Leaf mechanical properties in sclerophyll woodland and shrubland on contrasting soils. *Plant Soil* 276:95–113
- Savoie JM, Gourbiere F (1989) Decomposition of cellulose by the species of the fungal succession degrading *Abies alba* needles. *FEMS Microbiol Ecol* 62:307–314
- Schmidt O, Curry JP (1999) Effects of earthworms on biomass production, nitrogen allocation and nitrogen transfer in wheat–clover intercropping model systems. *Plant Soil* 214:187–198
- Scheu S, Theenhaus A, Jones TH (1999) Links between the detritivore and herbivore system: effects of earthworms and collembola on plant growth and aphid development. *Oecologia* 119:541–551
- Spain AV, Hutson BR (1983) Dynamics and fauna of the litter layers. In: Soils: an Australian Viewpoint. Division of Soils, CSIRO. CSIRO Publishing, Melbourne, pp 611–628
- Stock WD, Wienand KT, Baker AC (1995) Impacts of invading N₂-fixing *Acacia* species on patterns of nutrient cycling in two Cape ecosystems: evidence from soil incubation studies and ¹⁵N natural abundance values. *Oecologia* 101:375–382
- Thrall PH, Burdon JJ, Woods MJ (2000) Variation in the effectiveness of symbiotic associations between native rhizobia and temperate Australian legumes: interactions within and between genera. *J Appl Ecol* 37:52–65
- Thrall PH, Millsom DA, Jeavons AC, Waayers M, Harvey GR, Bagnall DJ, Brockwell J (2005) Seed inoculation with effective root-nodule bacteria enhances revegetation success. *J Appl Ecol* 42:740–751
- Troelstra SR, Wagenaar R, Smant W, Peters BAM (2001) Interpretation of bioassays in the study of interactions between soil organisms and plants: involvement of nutrient factors. *New Phytol* 150:697–706
- Tumbull MH, Goodall R, Stewart GR (1995) The impact of mycorrhizal colonization upon nitrogen source utilization and metabolism in seedlings of *Eucalyptus grandis* Hill ex Maiden and *Eucalyptus maculata* Hook. *Plant Cell Environ* 18:1386–1394
- Tumbull MH, Schmidt S, Erskine PD, Richards S, Stewart GR (1996) Root adaptation and nitrogen source acquisition in natural ecosystems. *Tree Physiol* 16:941–948
- Vesk PA, Mac Nally R (2006) The clock is ticking—revegetation and habitat for birds and arboreal mammals in rural landscapes of southern Australia. *Agric Ecosyst Environ* 112:356–366

- Vishnivetskaya T, Kathariou S, McGrath J, Gilichinsky D, Tiedje JM (2000) Low-temperature recovery strategies for the isolation of bacteria from ancient permafrost sediments. *Extremophiles* 4:165–173
- Walker V, Palmer GR, Voordouw G (2006) Freeze-thaw tolerance and clues to the winter survival of a soil community. *Appl Environ Microbiol* 72:1784–1792
- Wardle DA (2002) *Communities and ecosystems: linking the aboveground and belowground components*. Princeton University Press, Princeton
- Wardle DA, Bardgett RD, Klironomos JK, Setälä H, van der Putten WH, Wall DH (2004) Ecological linkages between aboveground and belowground biota. *Science* 304:1629–1633
- Warren CR (2006) Potential organic and inorganic N uptake by six *Eucalyptus* species. *Funct Plant Biol* 33:653–660
- Wood TG (1974) Field investigations on the decomposition of leaves of *Eucalyptus delegatensis* in relation to environmental factors. *Pedobiologia* 14:343–371
- Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J, Chapin T, Cornelissen JHC, Diemer M, Flexas J, Garnier E, Groom PK, Gulias J, Hikosaka K, Lamont BB, Lee T, Lee W, Lusk C, Midgley JJ, Navas ML, Niinemets U, Oleksyn J, Osada N, Poorter H, Poot P, Prior L, Pyankov VI, Roumet C, Thomas SC, Tjoelker MG, Veneklaas EJ, Villar R (2004) The worldwide leaf economics spectrum. *Nature* 428:821–827
- Yeates GW, Saggiar S, Hedley CB, Mercer CF (1999) Increase in ¹⁴C-carbon translocation to the soil microbial biomass when five species of plant-parasitic nematodes infect toots of white clover. *Nematology* 1:295–300